

Non-diagonal solutions to reflection equations in $su(n)$ spin chains

J. Abad and M. Rios

Departamento de Física Teórica, Facultad de Ciencias,
Universidad de Zaragoza, 50009 Zaragoza, Spain

Abstract

The reflection equations in a $su(3)$ spin chain with open boundary conditions are analyzed. We find non diagonal solutions to the boundary matrices. The corresponding hamiltonian is given. The solutions are generalized to $su(n)$.

The search for integrable models through the Yang-Baxter algebra has deserved considerable attention in the last years. For this purpose, quantum group representations give a systematic tool to obtain integrable systems. An aspect to take into account is which are the boundary conditions that those systems must verify in order to preserve integrability. In this line, the biggest development has been made for systems with periodic boundary conditions. So, the systems associated to the fundamental representations of the algebras $A_{(r)}$ can be found in ref. [1] and references therein. Other models associated to the algebra $A_{(1)}$ in nilpotent representations can be found in ref. [2]. Actually there is a lot of literature about models associated to other types of algebras.

An other aspect of the problem is to consider systems with open boundary conditions preserving integrability [3],[4]. The theoretical framework of the problem was set up in ref. [4] where boundary matrices are introduced and the relations they must fulfill in systems invariant under parity P , time reversal T and crossing unitarity are obtained, this formalism is extended the refs. [5]–[8] to systems symmetric under PT and non-invariant under crossing unitarity. A careful analysis in the framework of the algebraic structures was carry out in refs. [9] and [10]. The general solution for the boundary matrices in the models XXX , XXZ and XYZ can be found in [11]–[12]. Also diagonal solutions for the models $A_{(r)}$, $r \geq 2$ in their fundamental representation are obtained in [13] .

In this paper, we show non-diagonal solutions for the boundary matrices in models based on the algebras $A_{(r)}$, $r \geq 2$. To find them, we must solve a set of equations whose number increases exponentially with r , although the number of resulting incompatibilities that appears, which leading to simplifications, increases too.

As is well known, to find integrable systems with open boundary conditions, we must build the doubled transition operator [4], given by

$$U_{a,b}(\theta) = T_{a,c}(\theta) K_{c,d}^+(\theta) T_{d,b}^{-1}(-\theta) \quad (1)$$

with

$$T_{a,c}(\theta) = L_{a,a_1} \otimes L_{a_1,a_2} \otimes \cdots \otimes L_{a_{N-1},c}. \quad (2)$$

A graphic representation of the L operator is given in fig. 1. Tensorial product in the site spaces (vertical indices) and the usual product of matrices in the auxiliary space (horizontal indices) is understood .




Fig. 1

The $K^+(\theta)$ matrix, which determines the boundary condition must verify the well known reflection condition expressed by

$$R(\theta - \theta') [K^+(\theta) \otimes I] R(\theta + \theta') [K^+(\theta') \otimes I] = [K^+(\theta') \otimes I] R(\theta + \theta') [K^+(\theta) \otimes I] R(\theta - \theta'). \quad (3)$$

This relation is graphically expressed in fig. 2.




Fig. 2

Under this conditions, the operator

$$t(\theta) = K_{a,b}^-(\theta) T_{a,c}(\theta) K_{c,d}^+(\theta) T_{d,b}^{-1}(-\theta) \quad (4)$$

is a generator of constants of motion, since

$$[t(\theta), t(\theta')] = 0, \quad (5)$$

and then, an open local hamiltonian, given by

$$h = \left. \frac{dt(\theta)}{d\theta} \right|_{\theta=0}, \quad (6)$$

can be obtained.

If the system is invariant under parity, time reversal and crossing (case of $A_{(1)}$), then the K operator has the property

$$K^+(\theta) = K^{-t}(-\theta - \eta) \quad (7)$$

where t means transposition and η is the parameter of crossing property,

$$R_{a,c}^{b,d}(\theta) R_{c,e}^{d,f}(-\theta - 2\eta) \propto \delta_{a,e} \delta_{b,f}. \quad (8)$$

On the other hand, if the system is invariant under PT and not invariant under crossing symmetry and if there is a matrix M [5]–[7] such that

$$R_{a,b}^{c,d}(\theta) M_{b,e} R_{f,d}^{g,e}(-\theta - 2\eta) M_{f,h}^{-1} \propto \delta_{a,g} \delta_{c,h}. \quad (9)$$

then

$$K^+(\theta) = K^{-t}(-\theta - \eta) M^{-1}. \quad (10)$$

General solutions for the boundary matrices have been found in the models XXX , XXZ and XYZ [11], and diagonal solutions for the models of type $A_{(r)}$ with $r \geq 2$ diagonal solutions [13]. In this paper we present non diagonal solutions for $su(3)$ and $su(4)$ models, that we generalize afterwards to $su(n)$.

The matrix R is given [14] by

$$\begin{aligned} R(\theta) = & \sinh\left(\frac{n}{2}\theta + \gamma\right) \sum_{i=1}^n e_{i,i} \otimes e_{i,i} + \sinh\left(\frac{n}{2}\theta\right) \sum_{\substack{i,j=1 \\ i \neq j}}^n e_{i,j} \otimes e_{j,i} \\ & + \sinh(\gamma) \sum_{\substack{i,j=1 \\ i \neq j}}^n \exp\left((j-i - \frac{n}{2} \text{sign}(j-i))\theta\right) e_{i,i} \otimes e_{j,j}, \end{aligned} \quad (11)$$

that in case $n = 3$ becomes

$$R(\theta) = \begin{pmatrix} a & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & d & 0 & b & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & c & 0 & 0 & 0 & b & 0 & 0 \\ 0 & b & 0 & c & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & a & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & d & 0 & b & 0 \\ 0 & 0 & b & 0 & 0 & 0 & d & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & b & 0 & c & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & a \end{pmatrix}, \quad (12)$$

where

$$a(\theta) = \sinh\left(\frac{3}{2}\theta + \gamma\right), \quad (13a)$$

$$b(\theta) = \sinh\left(\frac{3}{2}\theta\right), \quad (13b)$$

$$c(\theta) = \sinh(\gamma)e^{\frac{\theta}{2}}, \quad (13c)$$

$$d(\theta) = \sinh(\gamma)e^{\frac{-\theta}{2}}. \quad (13d)$$

This model is PT invariant but not invariant under crossing symmetry, besides this, it verifies the property (9) with M equal to the identity I and $\eta = \gamma$.

We look for solutions to equation (3) with $K^+(\theta)$ of the form

$$K^+(\theta) = \begin{pmatrix} x_1(\theta) & y_1(\theta) & z_1(\theta) \\ x_2(\theta) & y_2(\theta) & z_2(\theta) \\ x_3(\theta) & y_3(\theta) & z_3(\theta) \end{pmatrix} \quad (14)$$

and fulfilling the condition

$$K^+(0) = I \quad (15)$$

As stated before, we are interested in non-diagonal solutions for $K^+(\theta)$. The resulting set of equations has not solution with all elements out of the diagonal in (14) different from zero. Those equations, instead, are satisfied when only one non-diagonal element and its symmetric are allowed to be different from zero, for instance,

$$K^+(\theta) = \begin{pmatrix} x_1(\theta) & 0 & z_1(\theta) \\ 0 & y_2(\theta) & 0 \\ x_3(\theta) & 0 & z_3(\theta) \end{pmatrix} \quad (16)$$

The solution that we find is

$$x_1(\theta) = K_{a+} \sinh\left(\epsilon_+ - \frac{3}{2}\theta\right) \quad (17a)$$

$$z_1(\theta) = K_{d+} e^{\frac{\theta}{2}} \sinh(3\theta) \quad (17b)$$

$$y_2(\theta) = K_{a+} e^{-\theta} \sinh\left(\epsilon_+ + \frac{3}{2}\theta\right) + K_{b+} e^{\frac{\theta}{2}} \sinh(3\theta) \quad (17c)$$

$$x_3(\theta) = K_{c+} e^{\frac{\theta}{2}} \sinh(3\theta) \quad (17d)$$

$$z_3(\theta) = K_{a+} e^{\theta} \sinh\left(\epsilon_+ + \frac{3}{2}\theta\right) \quad (17e)$$

where the K 's must be verified the relations

$$K_{c+} K_{d+} = K_{b+} (K_{b+} + K_{a+} e^{-\epsilon_+}). \quad (18)$$

Of course, if we impose $K_{b+} = 0$, we obtain the diagonal solutions [11], [13].

With this solutions, we can obtain through (10) the expression for the matrix $K^-(\theta)$ and then the hamiltonian (6). If we redefine the hamiltonian by subtracting a constant times the identity, in order to eliminate constant terms proportional to the length of the chain, it becomes

$$\begin{aligned} H^{su(3)} = & \frac{1}{2} \text{trace} (K^-(0)) \sum_{n=1}^{N-1} \left(J_9 (\lambda_n^3 \lambda_{n+1}^8 - \lambda_n^8 \lambda_{n+1}^3) + \sum_{\alpha=1}^8 (J_\alpha \lambda_n^\alpha \lambda_n^\alpha) \right) + \\ & + J_{1,3} \lambda_1^3 + J_{1,8} \lambda_1^8 + J_{1,4} \lambda_1^4 + J_{1,5} \lambda_1^5 + \\ & + \text{trace} (K^-(0)) (J_{N,3} \lambda_N^3 + J_{N,8} \lambda_N^8 + J_{N,4} \lambda_N^4 + J_{N,5} \lambda_N^5) \end{aligned} \quad (19)$$

with

$$J_1 = J_2 = J_4 = J_5 = J_6 = J_7 = 1 \quad (20a)$$

$$J_3 = J_8 = \cosh \gamma \quad (20b)$$

$$J_9 = \frac{\sinh(\gamma)}{\sqrt{3}} \quad (20c)$$

$$J_{1,3} = \frac{e^{-\frac{3\gamma}{2}} \sinh(3\gamma)}{3\gamma} \left(K_{b-}(1 + e^{2\epsilon_- + \gamma}) + \frac{K_{c-}K_{d-}}{K_{b-}}(e^{2\gamma} + e^{2\epsilon_- + \gamma}) \right) \quad (20d)$$

$$J_{1,8} = \frac{e^{-\frac{3\gamma}{2}} \sinh(3\gamma)}{\sqrt{3}\gamma} \left(-K_{b-}(1 + e^{2\epsilon_- + \gamma}) + \frac{K_{c-}K_{d-}}{K_{b-}} + e^{2\epsilon_- + \gamma} \right) \quad (20e)$$

$$J_{1,4} = \frac{-2e^{-\frac{\gamma}{2}} \sinh(3\gamma)}{\sinh \gamma} (K_{d-} + K_{c-}) \quad (20f)$$

$$J_{1,5} = \frac{-i2e^{-\frac{\gamma}{2}} \sinh(3\gamma)}{\sinh \gamma} (K_{d-} - K_{c-}) \quad (20g)$$

$$J_{N,3} = \frac{1}{\sinh \gamma} \left(K_{b+}(e^{2\epsilon_+} - 1) - \frac{K_{c+}K_{d+}}{K_{b+}}(2 + e^{2\epsilon_+}) \right) \quad (20h)$$

$$J_{N,8} = \frac{\sqrt{3}}{\sinh \gamma} \left(K_{b+}(e^{2\epsilon_+} + 1) - \frac{K_{c+}K_{d+}}{K_{b+}}e^{2\epsilon_+} \right) \quad (20i)$$

$$J_{N,4} = \frac{2}{\sinh \gamma} (K_{d+} + K_{c+}) \quad (20j)$$

$$J_{N,5} = \frac{2i}{\sinh \gamma} (K_{d+} - K_{c+}) \quad (20k)$$

and

$$\begin{aligned} \text{trace} (K^-(0)) = & \left(2 \cosh \gamma \sinh(\epsilon_- - \frac{3\gamma}{2}) + \sinh(\epsilon_- + \frac{3\gamma}{2}) \right) \left(\frac{K_{c-}K_{d-}}{K_{b-}} - K_{b-} \right) \\ & - K_{b-} e^{-\frac{\gamma}{2}} \sinh(3\gamma). \end{aligned} \quad (21)$$

In these expressions, we have removed the variables K_{a+} and K_{a-} by using (18) and a similar expression with the subindices (+) replaced by (-). So, the free parameters that we have in our solutions are

$$\epsilon_+, \epsilon_-, K_{b+}, K_{b-}, K_{c+}, K_{c-}, K_{d+}, K_{d-}.$$

The generalization to $su(n)$ follows from (17). The diagonal solution

$$K^+(\theta) = K_{i,i}^+ \delta_{i,j} \quad (22)$$

similar to that obtained in ref. [13], is

$$\begin{aligned} K_{j,j}^+(\theta) &= -\rho_+ e^{(2j+\frac{n}{2})\theta} \sinh(\epsilon_+ + \frac{n}{2}\theta) & \text{for } 1 \leq j \leq l_- \\ K_{l,l}^+(\theta) &= \rho_+ e^{(2l-\frac{n}{2})\theta} \sinh(-\epsilon_+ + \frac{n}{2}\theta) & \text{for } l_- + 1 \leq l \leq n \end{aligned} \quad (23)$$

being ρ and ϵ continuous parameters and l_- a discrete parameter with integer values $l_- \leq n-1$.

The non-diagonal solutions are

$$K_{1,1}^+(\theta) = \rho_{a+} \sinh(\epsilon_+ - \frac{n}{2}\theta), \quad (24a)$$

$$K_{n,n}^+(\theta) = \rho_{a+} e^{(n-2)\theta} \sinh(\epsilon_+ + \frac{n}{2}\theta), \quad (24b)$$

$$\begin{aligned} K_{j,j}^+(\theta) &= \rho_{a+} e^{(2j-n-2)\theta} \sinh(\epsilon_+ + \frac{n}{2}\theta) + \rho_{b+} e^{(2j-2-\frac{n}{2})\theta} \sinh(n\theta), \\ & j = 2 \cdots n-1, \end{aligned} \quad (24c)$$

$$K_{1,n}^+(\theta) = \rho_{d+} e^{(\frac{n}{2}-1)\theta} \sinh(n\theta), \quad (24e)$$

$$K_{n,1}^+(\theta) = \rho_{c+} e^{(\frac{n}{2}-1)\theta} \sinh(n\theta), \quad (24e)$$

the others components being zero. As before, the ρ 's parameters must verify the relation

$$\rho_{c+} \rho_{d+} = \rho_{b+} (\rho_{b+} + \rho_{a+} e^{-\epsilon_+}). \quad (25)$$

The solution (17) for $n=3$ is trivially obtained from (24) with the ρ 's parameters substituted by K 's ones.

As conclusions, we have shown a set of non-diagonal solutions for the boundary matrices in an open $su(n)$ spin chain. Others solutions, with the non zero elements out of the diagonal occupying others symmetric positions, can be obtained by means of similarity transformations.

In the analysis of the set of equations (3) and the solutions presented here, a heavy use of Mathematica computer program has been made.

Acknowledgements

We thank to professor J. Sesma for the careful reading of the manuscript and professor P. Kulish by useful comments. This work was partially supported by the Dirección General de Investigación Científica y Técnica, Grant No PB93-0302 and AEN94-0218

References

- [1] H.J. de Vega, Int. J. Mod. Phys. A4, (1989) 2371.
- [2] C. Gomez and G. Sierra, Phys. Lett. B285, (1992) 126.
- [3] V. Pasquier and H. Saleur, Nucl. Phys. B 330, (1990) 523.
- [4] E. K. Sklyanin, J. Phys. A21, (1988) 2375.
- [5] L. Mezincescu and R. I. Nepomechie, Int. J. Mod. Phys. A 6, (1991) 5281.
- [6] L. Mezincescu and R. I. Nepomechie, Int. J. Mod. Phys. A 7, (1992) 5657.
- [7] L. Mezincescu and R. I. Nepomechie, Jour. of Phys. A 24, (1991) L7.
- [8] N. Y. Reshetikhin and M. Semenov-Tian-Shansky, Lett. Math. Phys. 19, (1990) 133.
- [9] P. P. Kulish, R. Sasaki and C. Schwiebert, J. Math. Phys. 34, (1993) 286.
- [10] P. P. Kulish and E. K. Sklyanin, J. Phys. A25, (1992) 5963.
- [11] H. J. de Vega and A González-Ruiz, Boundary K-matrices for the XYZ , XXZ and XXX spin chains, preprint LPTHE-PAR 93-29 (hep-th/9306089).
- [12] T. Inami and H. Konno, J. Phys. A 27, (1994) L913.
- [13] H. J. de Vega and A González-Ruiz, Jour. of Phys. A 26, (1993) L519.
- [14] J. Abad and M. Rios, Integrable models associated to classical representations of $U_q(\widehat{sl(n)})$. University of Zaragoza preprint, DFTUZ 94/11.